

# How to use EpiPen® and EpiPen Jr® Auto-Injectors.

Remove the EpiPen® Auto-Injector from the carrier tube and follow these 2 simple steps:

---

## 1. Blue to the sky.



- Grasp with orange tip pointing downward
- Remove blue safety cap by pulling straight up – do not bend or twist

## 2. Orange to the thigh.



- Place the orange tip against the middle of the outer thigh
- Swing and push the auto-injector firmly into the thigh until it “clicks”
- Hold in place for 3 full seconds



Trusted for over 35 years.



After using EpiPen®, you must seek immediate medical attention or go to the emergency room. For the next 48 hours, you must stay close to a healthcare facility or be able to call 911.

# Be informed. Be prepared.



## Visit the [EpiPen.ca](https://www.EpiPen.ca) consumer site

- Learn about when and how to use your EpiPen® Auto-Injector
- Order **FREE** EpiPen® waistpacks, training kits and more
- Sign up for **FREE** EpiPen® expiration reminders and newsletters
- Browse the EpiPen® video gallery and FAQs

## Additional Resources

- Food Allergy Canada: [foodallergycanada.ca](https://www.foodallergycanada.ca)
- Allergy, Asthma & Immunology Society of Ontario: [allergyasthma.on.ca](https://www.allergyasthma.on.ca)
- Allergies Québec: [allergies-alimentaires.org](https://www.allergies-alimentaires.org)
- The Association of Allergists and Immunologists of Québec: [allerg.qc.ca](https://www.allerg.qc.ca)

EpiPen® and EpiPen Jr® (epinephrine) Auto-Injectors are indicated for the emergency treatment of anaphylactic reactions in patients who are determined to be at increased risk for anaphylaxis, including individuals with a history of anaphylactic reactions. Selection of the appropriate dosage strength is determined according to patient body weight. EpiPen® and EpiPen Jr® Auto-Injectors are designed as emergency supportive therapy only. They are not a replacement for subsequent medical or hospital care. After administration, patients should seek medical attention immediately or go to the emergency room. For the next 48 hours, patients must stay within close proximity to a healthcare facility or where they can call 911. To ensure this product is right for you, always read and follow the label. Please consult the Consumer Information leaflet in your product package for warnings and precautions, side effects, and complete dosing and administration instructions.

© 2021 Pfizer Canada ULC, Kirkland, Quebec H9J 2M5 • Toll free: 1-877-EPIPEN1 (1-877-374-7361)  
EpiPen®, EpiPen Jr® are registered trademarks of Mylan, Inc. licensed exclusively to its wholly-owned affiliate, Mylan Specialty, L.P.; sub-licensee, Pfizer Canada ULC, Kirkland, Quebec H9J 2M5 PP-EPI-CAN-0374-EN



Trusted for over 35 years.